

## Ranitidine Oral Solution

Products approved by Health Canada<sup>1</sup>:

Ingredient	Strength	Brand name	DIN	MFR
RANITIDINE	15MG/ML ORAL SOLUTION	NOVO-RANIDINE	02242940	NOP
RANITIDINE	15MG/ML ORAL SOLUTION	APO-RANITIDINE	02280833	APX

### Shortage management:

#### 1. Ensure valid indication for ranitidine solution.

- Acid suppressants are not recommended for uncomplicated gastroesophageal reflux in infants in the absence of warning signals such as poor weight gain, poor feeding, or unusual irritability.<sup>2,3</sup>

2. **If dose corresponds to tablet strengths** (75, 150, 300 mg), tablet can be crushed and mixed with soft food such as apple sauce immediately prior to administering.

#### 3. Prepare extemporaneous ranitidine liquid formulation<sup>4</sup>

- Ranitidine 15 mg/ml Simple Syrup Suspension<sup>5</sup>

<u>Ingredients</u>	<u>Quantity</u>
Ranitidine 150 mg	10 tablets
Distilled or Sterile water	50 mL
Simple syrup	qs to 100 mL

Directions:

1. In a mortar, crush tablets and triturate to a smooth powder.
2. Gradually add water. Mix well.
3. Pour into a graduated cylinder.
4. QS to final volume with simple syrup.
5. Transfer to final container and label.

Shake well. Pour immediately after shaking as suspension settles out rapidly.

Stability: 7 days at room temperature in amber plastic bottle.

- More recent investigations of the stability of extemporaneous ranitidine preparations in various strengths and vehicles, including Ora-Plus, report little or no loss of active ingredient for extended periods when stored at room temperature and refrigerated.<sup>6,7</sup> We do not have access to the original research papers and cannot assess the reliability of these results; furthermore, the form of ranitidine used (eg. powder, commercial

tablet, injectable, other) is unknown as are the specific procedures followed to compound. Therefore, we would recommend following the USP guidelines for a situation where the stability of an extemporaneous product is unknown i.e. **refrigerate and use within 14 days.**<sup>8</sup>

**Prepared by Karen Jensen, BSP, MSc. Reviewed by Carmen Bell, BSP  
medSask, March 2015.**

**References:**

1. Drug Product Database. Health Canada. Available at <http://webprod5.hc-sc.gc.ca/dpd-bdpp/index-eng.jsp>. Accessed 20Mar2015.
2. Lightdale JR, Gremse DA,. Gastroesophageal reflux: management guidance for the pediatrician. *Pediatrics*. 2013;131(5):e1684.
3. Mayo Clinic. Infant reflux – Treatment and drugs. Available at <http://www.mayoclinic.org/diseases-conditions/infant-acid-reflux/basics/treatment/con-20026253>. Accessed 20Mar2015.
4. Ranitidine. In: *Trissel's stability of compounded formulations*.(2012). Washington, DC: American Pharmacists Association.
5. Karnes HT, Harris SR, Garnett WR, et al. Concentration uniformity of extemporaneously prepared ranitidine suspension. *Am J Hosp Pharm*. 1989;46:304-7.
6. Ferreira MO, Bahia MF, Costa P. Stability of ranitidine hydrochloride in different aqueous solutions. *Eur J Hosp Pharm Sci*. 2004;10:60-3.
7. Lifshin LS, Fox JL. Stability of extemporaneously prepared ranitidine hydrochloride suspension. Paper presented at the ASHP Annual Meeting. 1992;120E.
8. Pharmaceutical Compounding – Nonsterile Preparations. U.S. Pharmacopeia. Available at [http://www.pharmacopeia.cn/v29240/usp29nf24s0\\_c795.html](http://www.pharmacopeia.cn/v29240/usp29nf24s0_c795.html). Accessed Feb 24, 2012.