

Pfizer-BioNTech


COMIRNATY™


mRNA COVID-19 vaccine

≥ 12 years

DIN 02527863

Grey Cap, Grey Border

 SUPPLIED:
multidose vial;
30 mcg/0.3 mL

 PREPARATION:
ready to use;
DO NOT DILUTE

 ADMINISTRATION:
IM injection

 APPROPRIATE
USE:

- 12 years & older
- **Primary series interval:** ≥ 21 days between doses 1 & 2; ≥ 28 days between doses 2 & 3 (immunocompromised)
- Additional considerations/Booster doses, eligibility, and intervals: see QR code



 BEYOND-USE
EXPIRY
DATE/TIME:

- **Unpunctured:**
 - 2°C to 8°C: use vaccine within 10 weeks
 - 8°C to 25°C: puncture vials within 12 hours*
*Including thaw time, if applicable
- **Punctured:** 2°C to 25°C: use vaccine within 12 hours of first puncture

 HANDLING:

- DO NOT DILUTE
- before use, invert vial gently 10 times to mix

 APPEARANCE:

- prior to inversion, may contain white to off-white opaque amorphous particles
- after inversion, should appear as white to off-white suspension;
- NO particulate matter/discolouration

 DOSE:

PRIMARY SERIES
0.3 mL (30 mcg mRNA)

BOOSTER DOSE
0.3 mL (30 mcg mRNA)
specific eligibility criteria, intervals, and doses
are available through QR code

 VIAL
STORAGE:

- between 2°C and 25°C for no more than 12 hours after first puncture
- protect from light when stored; can be handled in room light conditions

Pfizer-BioNTech


COMIRNATY™


mRNA COVID-19 vaccine

≥ 12 years

DIN 02527863

Grey Cap, Grey Border

 SUPPLIED:
multidose vial;
30 mcg/0.3 mL

 PREPARATION:
ready to use;
DO NOT DILUTE

 ADMINISTRATION:
IM injection

 APPROPRIATE
USE:

- 12 years & older
- **Primary series interval:** ≥ 21 days between doses 1 & 2; ≥ 28 days between doses 2 & 3 (immunocompromised)
- Additional considerations/Booster doses, eligibility, and intervals: see QR code



 BEYOND-USE
EXPIRY
DATE/TIME:

- **Unpunctured:**
 - 2°C to 8°C: use vaccine within 10 weeks
 - 8°C to 25°C: puncture vials within 12 hours*
*Including thaw time, if applicable
- **Punctured:** 2°C to 25°C: use vaccine within 12 hours of first puncture

 HANDLING:

- DO NOT DILUTE
- before use, invert vial gently 10 times to mix

 APPEARANCE:

- prior to inversion, may contain white to off-white opaque amorphous particles
- after inversion, should appear as white to off-white suspension;
- NO particulate matter/discolouration

 DOSE:

PRIMARY SERIES
0.3 mL (30 mcg mRNA)

BOOSTER DOSE
0.3 mL (30 mcg mRNA)
specific eligibility criteria, intervals, and doses
are available through QR code

 VIAL
STORAGE:

- between 2°C and 25°C for no more than 12 hours after first puncture
- protect from light when stored; can be handled in room light conditions