

Pfizer-BioNTech

COMIRNATY™ (5–11 years)

mRNA COVID-19 vaccine



SUPPLIED:
multidose
vial



PREPARATION:
reconstitute with
1.3 mL 0.9% Sodium
Chloride Injection, USP



ADMINISTRATION:
IM injection



**APPROPRIATE
USE:**

- 5 to 11 years old
- **2nd Dose:** authorized interval is at least 21 days from first dose; see QR code for optimal and mixed vaccine interval considerations
- **Additional/booster doses:** specific eligibility criteria, intervals, and doses are available through QR code



**BEYOND-USE
EXPIRY
DATE/TIME:**

- use vial within 6 months of manufactured date printed on carton/vial
- **Unreconstituted:**
 - use vials within 10 weeks when stored between 2°C and 8°C
 - use vials within 12 hours when stored between 8°C and 25°C
- **Reconstituted:** use vials within 12 hours when stored between 2°C and 25°C



HANDLING:

- inversion to take place during reconstitution
- no need to invert or swirl before drawing doses



APPEARANCE:

- ensure reconstituted
- white to off-white suspension, may contain opaque amorphous particles
- NO particulate matter/discolouration



DOSE:

PRIMARY SERIES
0.2 mL (10 mcg mRNA)

BOOSTER DOSE

To be determined

specific eligibility criteria, intervals, and doses are available through QR code



**VIAL
STORAGE:**

- between 2°C and 25°C for no more than 12 hours after reconstitution
- protect from light

REFERENCES:

- Comirnaty Product Monograph, BioNTech Manufacturing GmbH. Distributed by Pfizer Canada ULC. Kirkland QC. H9J 2M5. September 16, 2021. Available from <https://covid-vaccine.canada.ca/info/pdf/pfizer-biontech-covid-19-vaccine-pm1-en.pdf>

Pfizer-BioNTech

COMIRNATY™ (5–11 years)

mRNA COVID-19 vaccine



SUPPLIED:
multidose
vial



PREPARATION:
reconstitute with
1.3 mL 0.9% Sodium
Chloride Injection, USP



ADMINISTRATION:
IM injection



**APPROPRIATE
USE:**

- 5 to 11 years old
- **2nd Dose:** authorized interval is at least 21 days from first dose; see QR code for optimal and mixed vaccine interval considerations
- **Additional/booster doses:** specific eligibility criteria, intervals, and doses are available through QR code



**BEYOND-USE
EXPIRY
DATE/TIME:**

- use vial within 6 months of manufactured date printed on carton/vial
- **Unreconstituted:**
 - use vials within 10 weeks when stored between 2°C and 8°C
 - use vials within 12 hours when stored between 8°C and 25°C
- **Reconstituted:** use vials within 12 hours when stored between 2°C and 25°C



HANDLING:

- inversion to take place during reconstitution
- no need to invert or swirl before drawing doses



APPEARANCE:

- ensure reconstituted
- white to off-white suspension, may contain opaque amorphous particles
- NO particulate matter/discolouration



DOSE:

PRIMARY SERIES
0.2 mL (10 mcg mRNA)

BOOSTER DOSE

To be determined

specific eligibility criteria, intervals, and doses are available through QR code



**VIAL
STORAGE:**

- between 2°C and 25°C for no more than 12 hours after reconstitution
- protect from light

REFERENCES:

- Comirnaty Product Monograph, BioNTech Manufacturing GmbH. Distributed by Pfizer Canada ULC. Kirkland QC. H9J 2M5. September 16, 2021. Available from <https://covid-vaccine.canada.ca/info/pdf/pfizer-biontech-covid-19-vaccine-pm1-en.pdf>