

Pfizer-BioNTech


# COMIRNATY® ORIGINAL & OMICRON BA.4/BA.5


mRNA COVID-19 vaccine **bivalent**

≥ 5 years to  
< 12 years

DIN 02533197

Orange Cap,  
Orange Border

 SUPPLIED:  
multidose vial;  
**10 mcg\*/0.2 mL**

 PREPARATION:  
**DILUTE with 1.3 mL**  
0.9% Sodium Chloride  
Injection, USP

 ADMINISTRATION:  
IM injection

 APPROPRIATE  
USE:

- ≥ 5 years to < 12 years
- eligibility, intervals, and additional considerations: see QR code



 BEYOND-USE  
EXPIRY  
DATE/TIME:

- **Undiluted:**
  - 2°C to 8°C: use vaccine within 10 weeks
  - 8°C to 25°C: dilute vaccine within 12 hours\*
- \* Vials can be returned to fridge; however, any time kept above 8°C needs to be tracked and cannot exceed 12 hours.
- **Diluted:** 2°C to 25°C: use vaccine within 12 hours of dilution

 HANDLING:

- inversion to take place before and after dilution
- no need to invert or swirl before drawing doses
- do not shake

 APPEARANCE:

- ensure product has been diluted
  - white to off-white suspension
  - NO particulate matter/discolouration

 DOSE:

PRIMARY SERIES	BOOSTER DOSE
Specific eligibility criteria and intervals are available through QR code	
<b>0.2 mL</b> (10 mcg*)	<b>0.2 mL</b> (10 mcg*)

 VIAL  
STORAGE:

- between 2°C and 25°C for no more than 12 hours after dilution
- protect from light when stored; can be handled in room light conditions

\* Contains 5 mcg original & 5 mcg Omicron BA.4/BA.5 variant per 0.2 mL

Pfizer-BioNTech


# COMIRNATY® ORIGINAL & OMICRON BA.4/BA.5


mRNA COVID-19 vaccine **bivalent**

≥ 5 years to  
< 12 years

DIN 02533197

Orange Cap,  
Orange Border

 SUPPLIED:  
multidose vial;  
**10 mcg\*/0.2 mL**

 PREPARATION:  
**DILUTE with 1.3 mL**  
0.9% Sodium Chloride  
Injection, USP

 ADMINISTRATION:  
IM injection

 APPROPRIATE  
USE:

- ≥ 5 years to < 12 years
- eligibility, intervals, and additional considerations: see QR code



 BEYOND-USE  
EXPIRY  
DATE/TIME:

- **Undiluted:**
  - 2°C to 8°C: use vaccine within 10 weeks
  - 8°C to 25°C: dilute vaccine within 12 hours\*
- \* Vials can be returned to fridge; however, any time kept above 8°C needs to be tracked and cannot exceed 12 hours.
- **Diluted:** 2°C to 25°C: use vaccine within 12 hours of dilution

 HANDLING:

- inversion to take place before and after dilution
- no need to invert or swirl before drawing doses
- do not shake

 APPEARANCE:

- ensure product has been diluted
  - white to off-white suspension
  - NO particulate matter/discolouration

 DOSE:

PRIMARY SERIES	BOOSTER DOSE
Specific eligibility criteria and intervals are available through QR code	
<b>0.2 mL</b> (10 mcg*)	<b>0.2 mL</b> (10 mcg*)

 VIAL  
STORAGE:

- between 2°C and 25°C for no more than 12 hours after dilution
- protect from light when stored; can be handled in room light conditions

\* Contains 5 mcg original & 5 mcg Omicron BA.4/BA.5 variant per 0.2 mL