Pfizer-BioNTech COMIRNATY® Omicron XBB.1.5

mRNA COVID-19 vaccine

 \geq 12 years

DIN 02541823

Grey Cap, Grey Border



SUPPLIED:
multidose vial;
30 mcq/0.3 mL



PREPARATION: ready-to-use; DO NOT DILUTE





APPROPRIATE USE:

- · 12 years & older
- Eligibility, intervals, and additional considerations: see OR code





BEYOND-USE EXPIRY DATE/TIME:

- Check date on label (expiry date)
- · Unpunctured:
 - 2°C to 8°C: use vaccine within 10 weeks
 - 8°C to 25°C: puncture vial within 12 hours*
 - * Vials can be returned to fridge; however, any time kept above 8°C needs to be tracked and cannot exceed 12 hours.
- Punctured: 2°C to 25°C: use vaccine within 12 hours of first puncture



HANDLING

- DO NOT DILUTE
- · Before use, invert vial gently 10 times to mix; do not shake
- Can be handled in room light conditions



APPEARANCE:

- Prior to inversion, may contain white to off-white opaque amorphous particles
- After inversion, should appear as white to off-white suspension
- NO particulate matter/discolouration



0.3 mL (30 mcg)

Number of doses and intervals vary based on previous vaccination, immune status, and living situation. See QR code for details.



REFERENCE:

 Comirnaty Omicron X B8.1.5. Product Monograph. BioNTech Manufacturing GmbH. Distributed by Pfizer Canada ULC; 2023. Accessed October 2, 2023. https://covid-vaccine.canada.ca/info/ pdf/comirnaty-omicron-xbb-1-5-pm-en.pdf Financial contribution from



Agence de la santé a publique du Canada

12 OCT 2023 ©2023 medSask // Townes & Park Designs

Pfizer-BioNTech COMIRNATY® Omicron XBB.1.5

mRNA COVID-19 vaccine

 \geq 12 years

DIN 02541823

Grey Cap, Grey Border



SUPPLIED:

multidose vial; **30 mcg/0.3 mL**



PREPARATION: ready-to-use;

DO NOT DILUTE





APPROPRIATE USE:

12 years & older

 Eligibility, intervals, and additional considerations: see OR code





BEYOND-USE EXPIRY DATE/TIME:

- Check date on label (expiry date)
- Unpunctured:
- 2°C to 8°C: use vaccine within 10 weeks
 - 8°C to 25°C: puncture vial within 12 hours*
- *Vials can be returned to fridge; however, any time kept above 8°C needs to be tracked and cannot exceed 12 hours.
- Punctured: 2°C to 25°C: use vaccine within 12 hours of first puncture



HANDI ING.

- DO NOT DILUTE
- Before use, invert vial gently 10 times to mix; do not shake
- Can be handled in room light conditions



APPEARANCE:

- Prior to inversion, may contain white to off-white opaque amorphous particles
- After inversion, should appear as white to off-white suspension
- · NO particulate matter/discolouration



0.3 mL (30 mca)

Number of doses and intervals vary based on previous vaccination, immune status, and living situation. See QR code for details.



REFERENCE:

 Comirnaty Omicron X BB.1.5. Product Monograph. BioNTech Manufacturing GmbH. Distributed by Pfizer Canada ULC; 2023. Accessed October 2, 2023. https://covid-vaccine.canada.ca/info/ pdf/comirnaty-omicron-xbb-1-5-pm-en.pdf Financial contribution fro



olic Health Agence de la san ency of Canada publique du Cana

12 OCT 2023 ©2023 medSask // Townes & Park Designs