

AREXVY

non-live (recombinant) respiratory syncytial virus vaccine

RSV

≥ 60 years

DIN 02540207



SUPPLIED:

single-dose vial of antigen, vial of adjuvant suspension; needle/syringe not supplied



PREPARATION:

RECONSTITUTE prior to administration



ADMINISTRATION:

IM injection



APPROPRIATE USE:

- Not publicly funded in SK
- See QR code for recommendations for use.



ACIP



BEYOND-USE EXPIRY DATE/TIME:

- Check date on label (expiry date)
- **Not reconstituted:** 2°C to 8°C; use vaccine until expiry date.
- **Reconstituted:** 2°C to 25°C; use vaccine within 4 hours.



HANDLING:

- Gently swirl when reconstituting. DO NOT shake.



APPEARANCE:

- Reconstituted vaccine is opalescent, colourless to pale brownish liquid.
- NO foreign particulate matter/discolouration



DOSE:

- 0.5 mL



VACCINE SERIES:

- 1-dose

REFERENCES:

- [AREXVY product monograph](#)
- [Use of Respiratory Syncytial Virus Vaccines in Older Adults; Recommendations of the Advisory Committee on Immunization Practices 2023](#)

Financial contribution:



Public Health
Agency of Canada

Agence de la santé
publique du Canada