

IMOVAX[®] Polio

non-live (inactivated) poliomyelitis vaccine

IPV

≥ 2 months[‡]

DIN 01959042



SUPPLIED:
prefilled syringe;
needle not supplied



PREPARATION:
ready-to-use;
DO NOT DILUTE



ADMINISTRATION:
SUBQ injection



APPROPRIATE USE:

- Booster doses are not publicly funded in SK.
- See QR code for recommendations for use.



CIG



BEYOND-USE EXPIRY DATE/TIME:

- Check date on label (expiry date)
- 2°C and 8°C: use vaccine until expiry date.



HANDLING:

- **SHAKE VIGOROUSLY** before administration to re-suspend.



APPEARANCE:

- Clear, colourless suspension
- NO foreign particulate matter/discolouration



DOSE:

- 0.5 mL



SERIES/SCHEDULE:

- **PRIMARY SERIES:** Polio-containing vaccine series is given as part of Saskatchewan routine immunizations in childhood.
- **BOOSTER:** Single lifetime booster dose recommended for individuals ≥ 18 years at continued risk. See QR code for details.

[‡] Pharmacists can immunize those 5 years and older.

REFERENCES:

- [IMOVAX[®] Polio product monograph](#)
- [Saskatchewan Immunization Manual Chapter 10](#)
- [Canadian Immunization Manual: Poliomyelitis \(polio\) vaccines](#)
- [CATMAT Statement on Poliovirus and the International Traveller](#)

Financial contribution:



Public Health
Agency of Canada

Agence de la santé
publique du Canada

25 MAR 2024

©2024 medSask // Townes & Park Designs