

# IMOVAX<sup>®</sup> Rabies

non-live (inactivated) Human Diploid Cell Rabies vaccine

HDCV

Birth onwards<sup>‡</sup>

DIN 01908286



**SUPPLIED:**  
single-dose vial of lyophilized antigen, vial of adjuvant suspension for reconstitution; needle/syringe not supplied



**PREPARATION:**  
**RECONSTITUTE** prior to administration



**ADMINISTRATION:**  
IM injection



**APPROPRIATE USE:**

- Pre-exposure immunization is NOT publicly funded in SK.
- See QR code for recommendations for use.



CIG



**BEYOND-USE EXPIRY DATE/TIME:**

- Check date on label (expiry date)
- **Not reconstituted:** 2°C to 8°C; use vaccine until expiry date.
- **Reconstituted:** Use vaccine immediately.



**HANDLING:**

- Gently swirl when reconstituting. DO NOT shake.



**APPEARANCE:**

- Reconstituted vaccine is clear or slightly opalescent red to purplish red.
- NO particulate matter/change in physical appearance



**DOSE:**

• 1 mL



**SERIES/SCHEDULE:**

- **PRIMARY SERIES (PRE-EXPOSURE PROPHYLAXIS):**  
3-dose: 0, 7 days, 21 to 28 days\*
- **BOOSTER:** Recommended for some individuals with ongoing high risk of exposure to rabies. See CIG for details.

<sup>‡</sup> Pharmacists can immunize those 5 years and older.

\* as per the CIG

REFERENCES:

- [IMOVAX<sup>®</sup> Rabies product monograph](#)
- [Canadian Immunization Guide: Rabies vaccines](#)
- [CATMAT Statement on Travellers and Rabies Vaccine](#)

Financial contribution:



Public Health  
Agency of Canada

Agence de la santé  
publique du Canada

25 MAR 2024

©2024 medSask // Townes & Park Designs