

# IXIARO®

non-live (inactivated) Japanese  
encephalitis vaccine

JE

≥ 2 months<sup>‡</sup>

DIN 02333279



**SUPPLIED:**  
prefilled syringe;  
needle may be supplied



**PREPARATION:**  
ready-to-use;  
**DO NOT DILUTE**



**ADMINISTRATION:**  
IM injection



**APPROPRIATE  
USE:**

- Not publicly funded in SK
- See QR code for CIG recommendations for use.



CIG



**BEYOND-USE  
EXPIRY  
DATE/TIME:**

- Check date on label (expiry date)
- 2°C to 8°C: use vaccine until expiry date.



**HANDLING:**

- **SHAKE WELL** before administration to re-suspend.



**APPEARANCE:**

- White, cloudy liquid/suspension
- NO particulate matter/discolouration



**DOSE:**

- 0.5 mL



**SERIES/SCHEDULE:**

**PRIMARY SERIES:**

≥ 3 years<sup>‡</sup>: **Standard:** 2-dose: 0, 28 days

18 to 65 years: **Rapid/Accelerated:** 2-dose: 0, 7 days

**BOOSTER:**

May be required in some situations. See QR code for details.

<sup>‡</sup> Pharmacists can immunize those 5 years and older.

**REFERENCES:**

- [IXIARO® product monograph](#)
- [Canadian Immunization Guide: Japanese encephalitis vaccine](#)
- [CATMAT Statement on prevention of Japanese encephalitis](#)

Financial contribution:



Public Health  
Agency of Canada

Agence de la santé  
publique du Canada

25 MAR 2024

©2024 medSask // Townes & Park Designs