

M-M-R® II

live attenuated measles,
mumps and rubella virus vaccine

MMR

≥ 1 year[‡]

DIN 00466085



SUPPLIED:

single-dose vial of lyophilized powder, vial of sterile water diluent; needle/syringe not supplied



PREPARATION:

RECONSTITUTE
prior to
administration



ADMINISTRATION:

SUBQ
injection



APPROPRIATE USE:

- May be publicly funded in SK
- See QR code for recommendations for use

Note: Mumps and rubella included in all measles vaccines in Canada.



BEYOND-USE EXPIRY DATE/TIME: [^]

- Check date on label (expiry date)
- **Not reconstituted:** 2°C to 8°C (or in freezer above -50°C); use vaccine until expiry date.
- **Reconstituted:** 2°C to 8°C protected from light; use vaccine within 8 hours.

[^]PM includes excursion data.



CIG



SIM



HANDLING:

- Agitate to mix thoroughly when reconstituting.



APPEARANCE:

- Reconstituted vaccine is a clear yellow liquid with NO particulate matter/discolouration.
- Do not use if complete dissolution does not occur within 2 minutes.



DOSE:

- 0.5 mL



SERIES/SCHEDULE:

- 1 or 2 doses depending on age, previous immunization history, and risk factors: 0, ≥ 4 weeks. See CIG and SIM for details.

[‡] Pharmacists can immunize those 5 years and older.

REFERENCES:

- [M-M-R® II product monograph](#)
- [Saskatchewan Immunization Manual Chapter 10](#)
- [Canadian Immunization Guide: Measles vaccines](#)
- [Canadian Immunization Guide: Mumps vaccines](#)
- [Canadian Immunization Guide: Rubella vaccines](#)

Financial contribution:



Public Health
Agency of Canada

Agence de la santé
publique du Canada