

PREVNAR 20™

non-live (conjugate) pneumococcal vaccine

Pneu-C-20

≥ 6 weeks[‡]

DIN 02527049



SUPPLIED:
prefilled syringe;
needle not supplied



PREPARATION:
ready-to-use;
DO NOT DILUTE



ADMINISTRATION:
IM injection



APPROPRIATE USE:

- Not publicly funded in SK (Prevnar® 13, Pneumovax® 23 may be publicly funded)
- See QR code for adult recommendations for use.



CIG



BEYOND-USE EXPIRY DATE/TIME:[^]

- Check date on label (expiry date)
- 2°C and 8°C: use vaccine until expiry date.

[^] PM includes excursion data



HANDLING:

- **SHAKE VIGOROUSLY** before administration to re-suspend.



APPEARANCE:

- Homogeneous white suspension
- NO large particulate matter/discolouration



DOSE:

- 0.5 mL



SERIES/SCHEDULE:

- **Children/Adolescents:** SIM and CIG updates pending.
- **≥ 18 years:** 1-dose

[‡] Pharmacists can immunize those 5 years and older.

^{*} as per the CIG

REFERENCES:

- PREVNAR 20™ product monograph
- National Advisory Committee on Immunization (NACI) Public health level recommendations on the use of pneumococcal vaccines in adults, including the use of 15-valent and 20-valent conjugate vaccines (February 2023)
- National Advisory Committee on Immunization (NACI) Recommendations for public health programs on the use of pneumococcal vaccines in children, including the use of 15-valent and 20-valent conjugate vaccines (March 2024)

Financial contribution:



Public Health Agency of Canada

Agence de la santé publique du Canada