

PREVNAR 20™

non-live (conjugate) pneumococcal vaccine

Pneu-C-20

≥ 6 weeks[‡]

DIN 02527049



SUPPLIED:

prefilled syringe;
needle not supplied



PREPARATION:

ready-to-use;
DO NOT DILUTE



ADMINISTRATION:

IM injection



APPROPRIATE USE:

- May be publicly funded in SK
- See QR code for recommendations for use.



BEYOND-USE EXPIRY DATE/TIME:[^]

- Check date on label (expiry date)
- 2°C and 8°C: use vaccine until expiry date.

[^] PM includes excursion data



HANDLING:

- Store horizontally to minimize re-dispersion time.
- **SHAKE VIGOROUSLY** before administration to re-suspend.



CIG



SIM



APPEARANCE:

- Homogeneous white suspension
- NO large particulate matter/discolouration



DOSE:

- 0.5 mL



SERIES/SCHEDULE:

- ≥ 5 years: 1-dose. Timing depends on pneumococcal immunization history. See CIG and SIM for details.

[‡] Pharmacists can immunize those 5 years and older.

REFERENCES:

- [PREVNAR 20™ product monograph](#)
- [Saskatchewan Immunization Manual Chapter 10](#)
- [Canadian Immunization Guide: Pneumococcal vaccines](#)

Financial contribution:



Public Health
Agency of Canada

Agence de la santé
publique du Canada

21 JUNE 2024

©2024 medSask // Townes & Park Designs