

# RabAvert<sup>®</sup>

non-live (inactivated) Purified Chick Embryo Cell Rabies vaccine

PCECV

Birth onwards<sup>‡</sup>

DIN 02267667



**SUPPLIED:**  
single-dose vial of lyophilized powder, prefilled syringe containing sterile water diluent; two needles



**PREPARATION:**  
**RECONSTITUTE** prior to administration



**ADMINISTRATION:**  
IM injection



**APPROPRIATE USE:**

- Pre-exposure immunization is NOT publicly funded in SK.
- See QR code for recommendations for use.



CIG



**BEYOND-USE EXPIRY DATE/TIME:**

- Check date on label (expiry date)
- **Not reconstituted:** 2°C to 8°C; use vaccine until expiry date.
- **Reconstituted:** Use vaccine immediately.



**HANDLING:**

- Gently mix when reconstituting. DO NOT shake.



**APPEARANCE:**

- Reconstituted vaccine is a clear to slightly opalescent, colourless to slightly pink solution.
- NO foreign particulate matter/change in physical appearance



**DOSE:**

- 1 mL



**SERIES/SCHEDULE:**

- **PRIMARY SERIES (PRE-EXPOSURE PROPHYLAXIS):**  
3-dose: 0, 7 days, 21 to 28 days\*
- **BOOSTER:** Recommended for some individuals with ongoing high risk of exposure to rabies. See CIG for details.

<sup>‡</sup> Pharmacists can immunize those 5 years and older.

\* as per the CIG

REFERENCES:

- RabAvert<sup>™</sup> product monograph
- Canadian Immunization Guide: Rabies vaccines
- CATMAT Statement on Travellers and Rabies Vaccine

Financial contribution:



Public Health  
Agency of Canada

Agence de la santé  
publique du Canada

25 MAR 2024

©2024 medSask // Townes & Park Designs