

SHINGRIX

non-live (recombinant)
herpes zoster vaccine

RZV

≥ 50 years

Immunocompromised persons ≥ 18 years

DIN 02468425



SUPPLIED:

single-dose vial of lyophilized antigen, vial of adjuvant suspension for reconstitution; needle/syringe not supplied



PREPARATION:

RECONSTITUTE
prior to
administration



ADMINISTRATION:

IM injection



APPROPRIATE USE:

- Not publicly funded in SK
- See QR code for recommendations for use.



CIG



BEYOND-USE EXPIRY DATE/TIME:

- Check date on label (expiry date)
- **Not reconstituted:** 2°C to 8°C; use vaccine until expiry date.
- **Reconstituted:** 2°C to 8°C; use vaccine within 6 hours.



HANDLING:

- Shake vial gently when reconstituting.



APPEARANCE:

- Reconstituted vaccine is opalescent, colourless to pale brownish liquid.
- NO foreign particulate matter/discolouration



DOSE:

- 0.5 mL



SERIES/SCHEDULE:

- **2-dose:** 0, 2 to 6 months (0, 1 to 2 months if immunocompromised)

REFERENCES:

- [SHINGRIX product monograph](#)
- [Canadian Immunization Guide: Herpes zoster \(shingles\) vaccine](#)

Financial contribution:



Public Health
Agency of Canada

Agence de la santé
publique du Canada

25 MAR 2024

©2024 medSask // Townes & Park Designs