

TYPHIM Vi®

non-live (polysaccharide)
typhoid fever vaccine

Typh-I

≥2 years[‡]

DIN 02130955



SUPPLIED:
prefilled syringe;
choice of two needles



PREPARATION:
ready-to-use;
DO NOT DILUTE



ADMINISTRATION:
IM injection



**APPROPRIATE
USE:**

- Not publicly funded in SK
- See QR code for adult recommendations for use.



CIG



**BEYOND-USE
EXPIRY
DATE/TIME:**

- Check date on label (expiry date)
- 2°C to 8°C; use vaccine until expiry date.



HANDLING:

- **SHAKE WELL** before administration to re-suspend.



APPEARANCE:

- Clear, colourless solution
- NO foreign particulate matter/discolouration



DOSE:

- 0.5 mL



SERIES/SCHEDULE:

- **PRIMARY SERIES:** 1-dose
- **BOOSTER:** Recommended every 3 years for individuals at continued risk. See QR code for details.

[‡] Pharmacists can immunize those 5 years and older.

REFERENCES:

- TYPHIM Vi® product monograph
- Canadian Immunization Guide: Typhoid vaccines
- CATMAT Statement on International Travellers and Typhoid

Financial contribution:



Public Health
Agency of Canada

Agence de la santé
publique du Canada

25 MAR 2024

©2024 medSask // Townes & Park Designs