

YF-VAX®

live attenuated yellow fever virus vaccine

YF

≥ 9 months[‡]

DIN 00428833



SUPPLIED:

vial of lyophilized powder, vial of 0.9% Sodium Chloride diluent
YF vaccine is ordered directly from the manufacturer by designated YF vaccination centres ONLY.



PREPARATION:

RECONSTITUTE
prior to administration



ADMINISTRATION:

SUBQ
injection



APPROPRIATE USE:

- Not publicly funded in SK
- See QR code for adult recommendations for use.



CIG



BEYOND-USE EXPIRY DATE/TIME:

- Check date on label (expiry date)
- **Not reconstituted:** 2°C to 8°C: use vaccine until expiry date.
- **Reconstituted:** 2°C to 8°C; use vaccine within 60 minutes.



HANDLING:

- Gently swirl when reconstituting. DO NOT shake.
- Do NOT mix with other medicinal products, vaccines, or diluents.



APPEARANCE:

- Reconstituted vaccine is a slight pink-brown suspension.
- No foreign particulate matter/discolouration



DOSE:

- 0.5 mL



SERIES/SCHEDULE:

- **PRIMARY SERIES:** 1-dose
International Certificate of Vaccination or Prophylaxis (ICVP) must be provided to vaccine recipient; the ICVP becomes valid 10 days after immunization.
- **BOOSTER:** May be recommended 10 years after primary immunization in some individuals. See QR code for details.

[‡] Pharmacists can immunize those 5 years and older.

REFERENCES:

- YF-VAX® product monograph
- Canadian Immunization Guide: YF vaccine
- CATMAT Statement for Travellers and Yellow Fever

Financial contribution:



Public Health
Agency of Canada

Agence de la santé
publique du Canada

25 MAR 2024

©2024 medSask // Townes & Park Designs