

ABRYOVO™

non-live (subunit) respiratory
syncytial virus vaccine

RSV

- ≥ 60 years
- Pregnant individuals from 32 through 36 weeks gestational age

DIN 02544040



SUPPLIED:

single-dose vial of antigen,
prefilled syringe of diluent;
needle not supplied



PREPARATION:

RECONSTITUTE
prior to
administration



ADMINISTRATION:

IM injection



**APPROPRIATE
USE:**

- Not publicly funded in SK
- See QR code for recommendations for use.



ACIP



**BEYOND-USE
EXPIRY
DATE/TIME:**

- Check date on label (expiry date)
- **Not reconstituted:** 2°C to 8°C; use vaccine until expiry date.
- **Reconstituted:** 15°C to 30°C (do not refrigerate); use vaccine within 4 hours.



HANDLING:

- Gently swirl when reconstituting. DO NOT shake.



APPEARANCE:

- Reconstituted vaccine is a clear and colourless suspension.



DOSE:

- 0.5 mL



VACCINE SERIES:

- 1-dose

REFERENCES:

- [ABRYOVO™ product monograph](#)
- [Use of Respiratory Syncytial Virus Vaccines in Older Adults; Recommendations of the Advisory Committee on Immunization Practices 2023](#)

Financial contribution:



Public Health
Agency of Canada

Agence de la santé
publique du Canada