

Moderna

SPIKEVAX[®]

mRNA COVID-19 vaccine
Omicron KP.2 Formulation

≥ 6 months[‡]

DIN 02541270

Royal Blue Cap, Coral Blue Label



SUPPLIED:
multidose vial;
50 mcg/0.5 mL



PREPARATION:
ready-to-use
DO NOT DILUTE



ADMINISTRATION:
IM injection



APPROPRIATE USE:

- **6 months and older[‡]**
- Eligibility, intervals, and additional considerations: see QR code



BEYOND-USE EXPIRY DATE/TIME:

- **Check date on label** (expiry date) and **extensions** (in VDTs, Panorama, or CQE)
- **Unpunctured:**
 - 2°C to 8°C: use vaccine within 50 days
 - >8°C to 25°C: permitted exposure up to 12 hours* (not recommended storage)

* If the vial has been kept out of the fridge, it can be returned but the time kept above 8°C needs to be tracked and total time cannot exceed 12 hours.
- **Punctured:**
 - 2°C to 8°C: use vaccine within 24 hours* of first puncture
 - >8°C to 25°C: use vaccine within 12 hours* of first puncture

* Vials must be discarded 24 hours after first puncture when stored in the fridge. If the total time at room temperature (includes before and after puncture) exceeds 12 hours, the vials must be discarded.



HANDLING:

- Gently SWIRL before withdrawing each dose; **DO NOT SHAKE**
- Can be handled in room light conditions



APPEARANCE:

- White to off-white dispersion—may contain white or translucent particles
- **NO** foreign particulate matter/discolouration



DOSE:

AGE	DOSE [†]
≥ 6 months to < 5 years [‡]	0.25 mL (25 mcg)
≥ 5 years to < 12 years	0.25 mL (25 mcg)
≥ 12 years	0.5 mL (50 mcg)

[†] Number of doses and intervals vary based on previous vaccination, age, and immune status. See QR code for details.

[‡] Pharmacy professionals are not permitted to immunize individuals under 5 years of age.

REFERENCE:

- Spikevax Product monograph. Moderna Biopharma Canada Corp; 2024. Accessed September 17, 2024.
<https://covid-vaccine.canada.ca/info/pdf/spikevax-pm-en.pdf>

Financial contribution:



Public Health Agency of Canada
Agence de la santé publique du Canada